Modeling Potential Sites to Support Nearshore Marine Aquaculture on Hawai'i Island

Noelani Puniwai¹, Maria Haws¹, Barbara Gibson², James Potemra², Lisa Canale¹, Kohei Miyagi¹

¹University of Hawai'i at Hilo, Pacific Aquaculture and Coastal Resources Center ²University of Hawai'i at Mānoa, School of Ocean and Earth Science and Technology

Goal

Understand the potential for marine aquaculture along the Big Island's coasts

Aquaculture systems and models


- Caged aquaculture
 - brackish & oceanic finfish
- Line culture
 - algae & bivalves
- Intertidal Bottom Culture
 - bivalves
- Terrestrial Ponds
 - freshwater fish, shrimp, limu
 - ornamentals, pharmaceuticals
- Hawaiian Fishponds, ponds
 - range of fish species
 - limu, crustaceans
 - taro







Methods



1. Project Design & Area

- Focused on areas from shore to 3 Nautical Miles (State Waters)
 - GIS parameters: NAD 83 UTM Zone 5N & State Plan
- Sites identified by 100 hectare unique hexagons
 - 5,997 hexagons total ((±1.5 nm of 3 nm)
 - 3,227 hexagons (0-3 nm)
 - 1,578 hexagons (0-200m); 390,000 acres





2. Aquaculture Requirements

Biophysical Constraints

Water Conditions & Ocean Productivity	Ocean Conditions	Ocean Depth	Bottom Type
Salinity Turbidity Chlorophyll Temperature Chemical Factors (Oxygen & Pollution)	Tides Wave Height Flushing Roughness (Wind speed, Current Speed)	Depth Slope	Substrate Living Features

General Constraints

Current Zoning	Location & Economic Infrastructure	Physical Conflicts	Social Values
Marine Protected Areas Fishery Designated Areas Recreational Areas	Distance to Harbor Shoreline Access Shore-based Facility	Shipping lanes Buoys, Markers Military Dumping Area	Recreational Use Cultural Presence Viewshed

3-5. Data Gathering

- Technical Data Requirements

- Based on aquaculture Systems needs from literature & expert sources


According to Species needs from literature & expert sources

- GIS Data Compiled

- Over 75 different data layers compiled
- Layers used in the model included in the Geodatabase
- Metadata included with each layer

- Oceanographic Data (Satellite Imagery)

Ocean Currents, Wind Speed & Direction, Ocean color,
 Sea Surface Temperature, Ocean Tides, Ocean Waves




6. Scale & Extent of Data

- All datasets used must be available Islandwide
 - Site specific or incomplete coverage data not used
- Scale of data compared to size of hexagons and systems requirements
 - Important especially for satellite imagery

Wind Speed



October Chlorophyll minimum



7. Models

WHAT

WHY

HOW MUCH

Depth mbsynth50m

Water deep enough for water flow and little benthic impact.

> 4.5 m and < 30 m

SubstrateHabitat

Can't deploy equipment on coral, or near cliff faces

No Coral Reef / Precious Coral

Distance from Shore
Harbors/Ramps

About 30 min on a boat is the furthest you want to travel to access a site.

< 9 nm harbor or < 9 nm ramp


Salinity SGD & Streams

Known areas of freshwater streams and Submarine groundwater Discharge


50 m from streams and no SGD

8. Sample Model Results


- Red = Not good
- Yellow = Obstructions
- Green = Good



Habitat




Depth using Multibeam and 20' and 200' contours



Depth to Seafloor

Depth Zones

- Shallow (<20m) 772
- Medium (20-200m) 848
- Deep (>200m) 2304
- Adjacent to Land 580



Bathy" < 0 AND "Bathy" > - 31




Distance from Boat Ramp or Harbor

25 nautical miles from small boat harbors (SBH)


10 nautical miles from ramps, landings, SBH

Combined distances




Plankton – Chl-A concentrations

Oct mean (month of lowest values)
2003-2009 monthly values



Number of weeks Chl-A less than 0.05 mg/m³




Obstruction Factors

- No Buoys (100 m); 14
- No Cables (500 m); 26
- No Sewer Lines (100 m); 2
- No Lava Zone 1; 28
- No Marine Life Conserv Districts; 20
- No Offshore Installations; 10
- No Obstructions (100m); 9



Cautionary Factors (Socio-economic)

- Fishery Managed Areas
- Hawaiian Islands
 Humpback Whale NMS
- Ocean Rec. Design. Areas
- Public Parks & Beaches
- SCUBA Dive Sites
- Fish Aggregation Devices
- Body Surfing Sites



Caged Culture

b Habitat No Coral Presence 167 hex c Distance 25nm from harbor 1658 hex 10nm from a ramp 2810 hex	
b Habitat No Coral Presence 167 hex	
a Depth 20 - 200 m 777 hexago	S

b

Nearshore Aquaculture Models

128,000 acres

518 hex

Line Culture

	а	Depth	20 - 200 m	777 hexagons	
	b	Habitat	No Coral Presence	167 hex	D (
	С	Distance	25nm from harbor 10nm from a ramp	1658 hex 2810 hex	
	d	FreshH20	Perr Streams SGD	45 hex 226 hex	0
	C				
Ø	>	1			
-		a			
		S			
			b	d	36
				1/	

Intertidal Bottom Culture

-	_			
	а	Depth	4 - 30 m	772 hexagons
	b	Habitat	No Coral Presence	167 hex
١	С	Distance	25nm from harbor 10nm from a ramp	1658 hex 2810 hex
	d	FreshH20	Perr Streams SGD	45 hex 226 hex
1		ALC:		
1		a 🦠	С	3
		South States	5	~
	7		b	d

http://vs1b.geodata.hawaii.edu/aquaculture/cropmodel/



Results & Progress



3 Systems Modeled

- Inter-tidal Bottom Culture
- Line Culture
- Cage Culture

S. & N. Kohala

- Identified in all maps for potential
- Shallow Depth
- Close to Facilities

3 Systems Mapped

- Traditional fishponds
- Sites for Raised or Earthen Tanks
- Sites for Seaweed Culture

Depth


- Limited Shallow Areas
- MOORED SYSTEMS ONLY

Lack of Robust, Spatial Data




Questions to Think On...





Index to Site maps: for Hawaii, Kauai, Molokai, and Lanai



Site 2: Kiholo Bay, Kona Coast, Hawaii